



clinical investigations in critical care

Sodium Bicarbonate for the Treatment of Lactic Acidosis*

Sean M. Forsythe, MD; and Gregory A. Schmidt, MD, FCCP

Lactic acidosis often challenges the intensivist and is associated with a strikingly high mortality. Treatment involves discerning and correcting its underlying cause, ensuring adequate oxygen delivery to tissues, reducing oxygen demand through sedation and mechanical ventilation, and (most controversially) attempting to alkalinize the blood with IV sodium bicarbonate. Here we review the literature to answer the following questions: Is a low pH bad? Can sodium bicarbonate raise the pH *in vivo*? Does increasing the blood pH with sodium bicarbonate have any salutary effects? Does sodium bicarbonate have negative side effects? We find that the oft-cited rationale for bicarbonate use, that it might ameliorate the hemodynamic depression of metabolic acidemia, has been disproved convincingly. Further, given the lack of evidence supporting its use, we cannot condone bicarbonate administration for patients with lactic acidosis, regardless of the degree of acidemia. (CHEST 2000; 117:260–267)

Key words: acid-base; acidosis; alkalinizing therapy; bicarbonate; lactic acidosis; sodium bicarbonate

Abbreviations: DCA = dichloroacetate; DKA = diabetic ketoacidosis; SID = strong ion difference

Lactic acidosis, defined as a lactate concentration > 5 mmol/L and a pH < 7.35 , commonly complicates critical illness. Its causes are legion, including sepsis, cardiogenic shock, severe hypoxemia, hepatic failure, and intoxication. Many of these share reduced delivery of oxygen to cells or impaired use of oxygen in mitochondria, yet some are based in more complex derangements. For example, the lactic acidosis of sepsis is poorly understood but probably cannot be explained simply by tissue hypoxia, at least at the level of the whole body. Treatment of lactic acidosis involves discerning and correcting its underlying cause, ensuring adequate oxygen delivery to tissues, reducing oxygen demand through sedation and mechanical ventilation, and (in some ICUs) attempting to alkalinize the blood with IV sodium bicarbonate. Even in the face of maximal supportive therapy, lactic acidosis is associated with a mortality of 60 to 90%.^{1–4}

Although the use of sodium bicarbonate for the treatment of metabolic acidosis has been debated most heavily in the past 15 years, its use was questioned as far back as 1934.⁵ Nevertheless, it is still considered standard therapy and recommended in many textbooks and review articles,^{6–8} despite the lack of relevant clinical data supporting its effectiveness. Here we review the animal and human studies that bear on this complex, yet common, clinical conundrum.

We will not address the use of sodium bicarbonate in the bicarbonate-losing metabolic acidoses, such as those caused by diarrhea or renal tubular acidosis, in which replacement of lost bicarbonate is widely accepted. Although we touch on the use of bicarbonate in diabetic ketoacidosis (DKA) and some other forms of metabolic acidosis, our discussion and conclusions largely relate to the single most common cause of severe lactic acidosis in the modern ICU, that caused by severe sepsis.⁴ We find little evidence that sodium bicarbonate is detrimental in these conditions, but its use fails the most basic criteria expected of any drug—efficacy.

Those who continue to advocate the use of sodium bicarbonate for lactic acidosis generally use the following chain of reasoning (explicitly or implicitly):

*From the Department of Medicine (Dr. Forsythe) and the Section of Pulmonary and Critical Care (Dr. Schmidt), University of Chicago School of Medicine, Chicago, IL.

Manuscript received July 8, 1999; revision accepted July 15, 1999.

Correspondence to: Gregory A. Schmidt, MD, FCCP, Section of Pulmonary and Critical Care: MC6026, 5841 S. Maryland Ave, Chicago, IL 60637; e-mail: gschmidt@medicine.bsd.uchicago.edu

1. A low pH, in and of itself, is harmful (most notably by impairing cardiovascular function).
2. Sodium bicarbonate can increase the pH when infused IV.
3. Raising the pH with sodium bicarbonate improves cardiovascular function or some other relevant outcome.
4. Any adverse effects of sodium bicarbonate are outweighed by its benefits.

We address below each of these points in turn.

IS A LOW PH BAD?

Proteins, which underlie the function and structure of human cells, contain areas of both positive and negative charge and are thereby sensitive to the pH of the surrounding milieu. One can conceive of innumerable ways in which excess acid could impair protein function and, by extension, the function of the whole body. Yet it is overly simplistic to assume that the clinician's window on acid-base state, the arterial blood pH, reflects accurately the pH at a (likely more important) cellular level. For example, a 50% reduction in blood flow to a tissue causes the arterial-venous PCO₂ difference to double (as long as CO₂ production and excretion remain constant) as predicted by the Fick principle. This will substantially raise the tissue PCO₂ and lower the local intracellular pH. At the same time, neither the arterial pH nor PCO₂ changes at all, both failing completely to reveal the tissue acidosis. Further clouding the value of the arterial blood pH, there may be different acid-base states in different cells of a single organ, or within different organs of a single patient. Mitochondrial pH may be even more crucial than cellular (cytoplasmic) pH, as these organelles are the site of energy production. In experiments involving isolated rat hepatocytes, the mitochondrial membrane pH gradient did not change when extracellular pH was lowered from 7.40 to 6.9.⁹

Because adults with acidosis generally also have sepsis, hypoxemia, intoxication, or hypoperfusion, discerning the physiologic effects of low pH from those of endotoxemia, hypoxemia, and so on is a challenging task. In isolated animal heart muscle preparations in which the pH of the perfusate is lowered, acidosis generally reduces contractile function,¹⁰ sometimes severely.¹¹ Human ventricular muscle excised during open-heart surgery displays only modestly reduced contractility in the face of severe acidosis.¹² The cardiac depressant effect may be caused by inorganic phosphate-mediated impairment of actin-myosin crossbridge cycling, disruption of energy production,^{11,13} interference with calcium delivery to myofilaments, or decreased sensitivity of

contractile proteins to calcium.¹⁴ In whole-animal preparations, the effects of acidosis are more difficult to discern, because of competing effects of acidosis on contractility, heart rate, vascular tone, and the adrenal and sympathetic systems. Various investigators have controlled heart rate, preload, and afterload, finding that acidosis caused contractility to remain constant,¹⁵ decrease marginally,¹⁶ or transiently rise and then fall.¹⁷

The increasing experience with permissive hypercapnia for patients with ARDS or status asthmaticus, in which hypercapnia and acidemia are tolerated to avoid alveolar overdistention, has changed many clinicians' perspective about the adverse impact of acidemia. In sedated and ventilated patients with ARDS, rapid intentional hypoventilation (pH falling from 7.40 to 7.26 in 30 to 60 min) lowered systemic vascular resistance while cardiac output rose. Mean systemic arterial pressure and pulmonary vascular resistance were unchanged.¹⁸ Further, in many studies of patients undergoing permissive hypercapnia, a pH of well below 7.2 was tolerated well,¹⁸⁻²⁵ as it is in young patients with DKA,²⁶ children with "super-carbia,"²⁷ and those with grand mal seizures.²⁸⁻³⁰ The feared consequences of acidemia, projected from the experience with patients having lactic acidosis (and, usually, concomitant sepsis), failed to materialize. In normal subjects who rebreathed carbon dioxide, QT dispersion was increased, which could signal a risk of arrhythmia.²⁴ With data now available for many patients permissively hypoventilated, the systemic hemodynamic effects are quite small even as the pH falls to 7.15, with the typical patient experiencing no change or small increases in cardiac output and BP. Rhythm disturbances have not been a problem. Patients whose pH levels fall far below 7.0 are fewer in number, so firm conclusions cannot be drawn, but they similarly tolerate their acidemia. The current practice of permissive hypercapnia does not generally include an attempt to alkalinize the blood to compensate for respiratory acidosis.

Cardiac contractile response to catecholamines is also impaired by acidosis, perhaps mediated by a decline in β -receptors on the cell surface.^{13,31,32} Other potentially detrimental cardiovascular effects of acidosis include reduced resuscitability from induced ventricular fibrillation, which has been shown in rats but not in dogs or pigs,³³⁻³⁵ impaired load tolerance of the right ventricle,³⁶ and altered renal blood flow (both increased and decreased, depending on the degree of acidemia).³⁷ Diaphragmatic contractility is reduced also by respiratory acidosis,³⁸ but apparently not by metabolic acidosis.³⁹

Paradoxically, acidosis may have protective effects in critical illness. A low pH has been shown to delay

the onset of cell death in isolated hepatocytes exposed to anoxia⁹ and to chemical hypoxia.⁴⁰ Correcting the pH took away the protective effect and accelerated cell death. In addition, acidosis during reperfusion limits myocardial infarct size.^{41,42}

In summary, although a very low pH has negative inotropic effects in isolated hearts, the whole-body response in patients is much less clearly detrimental. Although clinical shock and metabolic acidosis often coincide, the striking discordance between the clinical course and outcome of patients with (usually septic) lactic acidosis compared with those who have DKA or ventilatory failure suggests that the low pH itself does not crucially underpin the hemodynamic collapse of these ill patients. Independent of the acidemia, the lactate ion may be significant, because lactate buffered to a pH of 7.4 can cause decreased cardiac contractility in animal models.⁴³

CAN SODIUM BICARBONATE RAISE THE pH IN VIVO?

It seems straightforward that adding a base to acidic blood will raise the pH—the reality is more complex. First, bicarbonate is not one of the independent determinants of the blood pH. Rather, these include the difference between the total concentrations of strong cations and anions (the strong ion difference, [SID]); the total concentration of weak acids; and the PaCO₂.⁴⁴ Administration of exogenous sodium bicarbonate increases the SID (which tends to raise the pH) because sodium is a strong cation and bicarbonate is not a strong ion at all, but at the same time it elevates the PaCO₂ (which tends to lower pH). In patients with lactic acidosis receiving mechanical ventilation, a modest infusion of sodium bicarbonate (2 mmol/kg for 15 min IV) boosted PaCO₂ from 35 to 40 mm Hg.²

Second, bicarbonate administration may engender metabolic reactions that may themselves alter (SID), the total concentration of weak acids, or PaCO₂. For example, in animals and humans, bicarbonate infusion can augment the production of lactic acid, a strong anion.^{45–50} Mechanisms to explain this remain speculative, but include a shift in the oxyhemoglobin-saturation relationship⁵¹; enhanced anaerobic glycolysis, perhaps mediated by the pH-sensitive, rate-limiting enzyme phosphofructokinase; and changes in hepatic blood flow or lactate uptake.⁵²

In animal models of lactic acidosis, sodium bicarbonate does not predictably raise the arterial pH. In some studies, pH remained constant or fell.^{46,47} Most whole-animal studies, however, have shown that pH can be raised and even normalized.^{16,48,49,53–55} Most relevant to clinical practice, in two studies of patients with lactic

acidosis receiving mechanical ventilation, IV infusion of sodium bicarbonate in dosages of 2 mmol/kg for 15 min or 1 mmol/kg for 1 to 2 min raised the pH only 0.14 and 0.05 units, respectively.^{2,56}

Yet, the body has multiple compartments separated by membranes of differing permeabilities and systems of active transport. Even when sodium bicarbonate added to the central veins reliably elevates the arterial pH, its effects on the cerebrospinal fluid and intracellular spaces may not be concordant. This could happen because carbon dioxide, produced when bicarbonate reacts with metabolic acids, diffuses readily across cell membranes and the blood-brain barrier, whereas bicarbonate cannot. Alternatively, as discussed above, bicarbonate may provoke reactions within cells, vessels, or organs that lower the local SID or raise the local PCO₂.

Sodium bicarbonate lowered cerebrospinal fluid pH in dogs with DKA,⁵⁷ dogs being resuscitated after a ventricular fibrillation arrest,⁵⁸ and two patients with DKA, in whom it was associated with a decrease in mental status.⁵⁹ Intracellular pH has been measured in cells or animals using nuclear magnetic resonance spectroscopy, pH-sensitive fluorescent dyes, intramuscular electrodes, and the distribution of carbon 14-labeled dimethadione (Table 1). The results are inconsistent, with intracellular pH rising in one study,⁵⁵ not changing in six studies,^{11,34,49,53,60,61} falling in nine studies,^{46,47,62–68} and either rising or falling depending on the buffer used in two investigations.^{69,70} In various studies, intracellular pH has been shown to fall with bicarbonate in RBCs,⁴⁶ muscle,⁶⁸ liver,⁴⁷ lymphocytes,⁶⁶ and brain.⁶³ We are aware of only one human study involving determination by magnetic resonance spectroscopy of intracellular pH in the brain.⁶⁵ When these normal volunteers were given sodium bicarbonate IV, brain pH fell significantly.

In summary, sodium bicarbonate can raise the blood pH when given IV. In contrast, this therapy fails to augment reliably the intracellular pH. Indeed, intracellular pH falls in most animal models and in most organs studied, but the effect is variable.

DOES INCREASING THE BLOOD pH WITH SODIUM BICARBONATE HAVE ANY SALUTARY EFFECTS?

The most direct question to pose regarding sodium bicarbonate therapy is whether it improves the problems that prompt its use. Namely, does it correct hemodynamics, “buy time” for other interventions, or improve outcome?

In isolated rat or rabbit hearts perfused with acidic solutions, bicarbonate fails to augment ventricular

Table 1—Results of Studies of the Effect of Bicarbonate on Intracellular pH*

Source	Subject	Acidosis	Method of Measuring Intracellular pH	Serum pH	Intracellular pH
Beech et al. ⁵⁵	Rat	DKA, shock	³¹ P NMR	↑	↑ (heart)
Rhee et al. ⁴⁹	Dog	Hypoxic lactic	³¹ P NMR	↔	↔ (heart)
Beech et al. ⁵³	Rat	Hypotensive lactic	C2 NMR	↑	↔ (muscle)
Bollaert et al. ⁶⁰	Rat	Septic (LPS)	³¹ P NMR	↑	↔ (muscle)
Shapiro ¹¹	Rat heart	Acidic perfusate	³¹ P NMR	↑	↔ (heart)
Thompson et al. ⁶¹	Rat	None	³¹ P NMR	↑	↔ or ↓ depending on route (IV or IP)
Kette et al. ³⁴	Pig	Cardiac arrest	Electrode	↑	↔ (heart)
Arieff et al. ⁴⁶	Dog	Phenformin lactic	¹⁴ C DMO	↔	↓ (liver, RBC)
Graf et al. ⁴⁷	Dog	Hypoxic lactic	¹⁴ C DMO	↔	↓ (liver)
Bersin and Arieff ⁶²	Dog	Hypoxic lactic	¹⁴ C DMO	↓	↓ (liver)
Shapiro et al. ⁶³	Rat	NH ₄ Cl, hypercapnic	³¹ P NMR	↑	↓ (brain)
Shapiro et al. ⁶⁴	Rat	NH ₄ Cl	³¹ P NMR	↑	↓ (liver)
Arieff et al. ⁶⁵	Animal	Phenformin lactic	Not stated	↑	↓ (liver and muscle)
Nakashima et al. ⁶⁵	Human	None	³¹ P NMR	↓	↓ (brain)
Bjerneroth et al. ⁶⁶	Lymphocytes	Acidic buffer	Fluorescent dye	↑	↓
Ritter et al. ⁶⁷	Platelets	Acidic buffer	Fluorescent dye		↓
Levrant et al. ⁷⁰	Rat hepatocytes	Acidic buffer	Fluorescent dye		↑ or ↓ depending on buffer
Goldsmith et al. ⁶⁹	Leukocytes	Acidic buffer	Fluorescent dye		↑ or ↓ depending on buffer

*NH₄Cl = ammonium chloride; NMR = nuclear magnetic resonance; DMO = dimethylxozolidine; IP = intraperitoneal; LPS = lipopolysaccharide; ↑ = increase; ↓ = decrease; ↔ = no change.

contractility.^{11,71} In whole animals (including various models of metabolic acidosis), the effects of bicarbonate on ventricular function are more difficult to tease out from its impact on systemic vessels. Further, one must take care not to interpret these studies too simplistically. For example, a fall in BP is not necessarily detrimental (if cardiac output rises). Nevertheless, these studies uniformly fail to reveal any hemodynamic benefit for sodium bicarbonate when compared with iso-osmolar saline solution.^{16,45–49,53,55,62,64,72–74} When it has been measured, cardiac output either does not change^{48,49,74} or falls.^{11,46,47} Right ventricular contractility^{72,74} and hepatic blood flow^{46,47} fall. Perhaps the most careful study of left ventricular function involved L-lactic acid infusion in anesthetized, ventilated, β-blocked, and atrially paced dogs. Before and after bicarbonate infusion, the left ventricular pressure-volume relationship was determined with a ventricular Millar catheter and three orthogonal pairs of ultrasonic crystals imbedded in the ventricular walls.¹⁶ Although sodium bicarbonate increased the arterial pH and did not increase lactate concentrations, mean arterial pressure fell, and cardiac output and ventricular contractility (slope of the end-systolic pressure-volume relationship) did not change. The hemodynamic effects were indistinguishable from those of saline solution.

There have been two studies of the hemodynamic impact of sodium bicarbonate in human lactic acidosis.^{2,56} In both studies, patients were receiving mechanical ventilation, had a mean blood lactate between 7 mmol/L and 8 mmol/L, and were receiving

continuous infusions of vasoactive drugs (except one patient in one study). Although sodium bicarbonate raised pH and serum bicarbonate concentrations, it did not improve hemodynamics or catecholamine responsiveness. Specifically, bicarbonate was indistinguishable from saline with regard to heart rate, central venous pressure, pulmonary artery pressure, mixed venous oxyhemoglobin saturation, systemic oxygen delivery, oxygen consumption, arterial BP, pulmonary artery occlusion (wedge) pressure, and cardiac output.^{2,56} These findings suggest that the commonly observed hemodynamic response to bicarbonate administration in patients treated with vasoactive drug infusions may simply be one of preload augmentation (rather than enhanced catecholamine responsiveness). When the most severely acidemic (pH range 6.9 to 7.2) subset of patients was analyzed separately, these negative findings persisted.² This result does not support the practice of some physicians who withhold bicarbonate from patients with mild acidemia but feel compelled to give it to those with acidemia of greater magnitude. Indeed, if there are negative effects of bicarbonate infusion, there are reasons to expect that this subset of patients will suffer disproportionately (*ie*, develop more profound paradoxical intracellular acidosis).

Outcome is difficult to measure because animal models have a nearly 100% mortality and human trials have generally not been designed to detect differences in survival or other (nonhemodynamic) measures of outcome. In both prospective and retrospective studies of patients with DKA treated with or without sodium bicarbonate, there were no dif-

ferences in the neurologic status, incidence of hypokalemia or hypoglycemia, or rate of correction of acidemia,^{75,76} but there was a suggestion of delayed clearance of ketones and lactate in patients given bicarbonate.^{77,78} Dichloroacetate (DCA) infusion, which, like sodium bicarbonate infusion, effectively raises serum pH in critically ill patients with lactic acidosis (but lowers lactate concentrations), also has no apparent hemodynamic benefit and does not improve survival.³

The only study that has shown any positive effect on outcome is in the setting of canine ventricular fibrillation.⁷⁹ Dogs resuscitated from prolonged arrest who were given bicarbonate had improved return of the circulation, less neurologic deficit, and greater survival to 24 h. On the other hand, several other studies in both humans and animals do not support these data. A study in dogs showed no effect of respiratory or metabolic acidosis on defibrillation threshold.³⁵ Sodium bicarbonate had no detectable impact on myocardial cell pH or resuscitability from ventricular fibrillation in pigs in one study³⁴ and worsened coronary perfusion pressure and resuscitability in yet another.⁸⁰ Its use was associated with poorer outcomes in a retrospective study of human cardiopulmonary arrest.⁸¹

In summary, no controlled study has shown improved hemodynamics attributable to sodium bicarbonate infusion, regardless of the effect on pH, and many show worsening of some hemodynamic variable. It is significant that such negative findings include two studies in critically ill humans receiving infused catecholamines, the subset of patients who might be expected to benefit most dramatically.

DOES SODIUM BICARBONATE HAVE NEGATIVE SIDE EFFECTS?

The most obvious side effects of sodium bicarbonate are the fluid and sodium load. This can cause hypervolemia, hyperosmolarity, and hypernatremia.⁸² Sodium bicarbonate given as a rapid IV bolus can cause a transient fall in mean arterial pressure and a transient rise in intracranial pressure⁸³ that is probably related to its hypertonicity, and this is alleviated when given as a slow IV infusion. Sodium bicarbonate has been shown in three studies to lower PaO₂ from 5 to 15 mm Hg in both acidemic animals and nonacidemic patients with congestive heart failure.^{57,84} The mechanism for this is unclear, but the authors speculated that there might be worsening of intrapulmonary shunt.

When normal human volunteers were made acidemic with acetazolamide and then corrected with sodium bicarbonate, the acute correction of the pH

caused increased hemoglobin affinity for oxygen that worsened oxygen delivery. This effect lasted about 8 h.⁵¹ This was thought to be caused by the immediate nature of the Bohr effect and the delayed nature of the 2,3-diphosphoglycerate effect on hemoglobin-oxygen affinity.

Lactate concentrations increased with sodium bicarbonate infusion (compared with control subjects) in animal studies of hypoxic lactic acidosis,^{45,47,49} phenformin-induced lactic acidosis,^{46,68} hemorrhagic shock,⁴⁸ and DKA.^{55,57} This finding has also been reported in cases of chronic lactic acidosis associated with malignancies.⁸⁵ It is possible that lactate rises in these settings because of impaired oxygen delivery to tissues. Even if enhanced lactate production does not signal cellular hypoxia, bicarbonate-induced hyperlactatemia may be detrimental inasmuch as lactate itself has potentially detrimental actions, as discussed earlier.

Serum ionized calcium concentration is reduced by sodium bicarbonate infusion. In a randomized, controlled study of ICU patients with lactic acidosis, sodium bicarbonate lowered ionized calcium from 0.95 to 0.87 mmol/L.² In an animal study of cardiac arrest, sodium bicarbonate decreased ionized calcium, although this had no apparent detrimental effects.⁷⁹ Because left ventricular contractility has been shown to vary directly with ionized calcium concentration,⁸⁶ any beneficial effects of pH correction may be masked by hypocalcemic ventricular depression.

A single study in patients with lactic acidosis treated with sodium bicarbonate failed to reveal any significant changes in venous lactate concentration, hemoglobin affinity for oxygen, total body oxygen consumption, oxygen extraction ratio, transcutaneous oxygen pressure, serum sodium concentration, or osmolality.⁵⁶ However, the dose of bicarbonate (1 mmol/kg) was small. Another study of three patients treated with sodium bicarbonate (mean dose, 90 mEq) during cardiac arrest revealed a rise in osmolality from 308 to 343 mosm/kg.⁵⁸

In summary, many potentially detrimental effects of bicarbonate administration have been identified, but their clinical relevance has not been established.

OTHER ALKALINIZING THERAPIES: CARBICARB, DICHLOROACETATE, TRIS-HYDROXYMETHYL AMINOMETHANE, AND DIALYSIS

Carbicarb is an equimolar mixture of sodium carbonate and sodium bicarbonate. Compared with sodium bicarbonate, Carbicarb raises the SID (thereby the pH) far more^{45,48,49,54,62} and boosts the PCO₂ far less^{48,49,64} when given IV to animals with

metabolic acidosis. To the extent that the failure of sodium bicarbonate to effect hemodynamic improvement is caused by the generation of carbon dioxide, Carbicarb might be superior. Carbicarb more consistently increases intracellular pH,^{49,62,64} and although it improved hemodynamics in two studies,^{11,49} it did not in three others.^{48,62,64}

DCA is a compound that probably works by stimulating the pyruvate dehydrogenase complex, the rate-limiting enzyme that regulates the entry of pyruvate into the tricarboxylic acid cycle, thus promoting the clearance of accumulated lactate. In addition, DCA increases myocardial glucose utilization and contractility.⁸⁷ Although several animal and clinical trials showed that DCA could raise pH and bicarbonate concentration while lowering lactate concentrations, with little apparent toxicity,^{52,88,89} a large, multicenter, placebo-controlled trial in patients with lactic acidosis failed to confirm improved hemodynamics or outcome.³ This drug is not available commercially. Tris-hydroxymethyl aminomethane is a weak alkali that penetrates cells easily. Its potential to raise blood and intracellular pH without producing carbon dioxide has been confirmed in animal models of metabolic acidosis. Tris-hydroxymethyl aminomethane also improved myocardial contraction and relaxation in an isolated rabbit heart preparation.⁷¹ Although tris-hydroxymethyl aminomethane is commercially available, complications of hyperkalemia, hypoglycemia, extravasation-related necrosis, and neonatal hepatic necrosis are likely to limit its use.

There have been many case studies that report the use of dialysis to control the volume and sodium loads that accompany sodium bicarbonate infusion. In most of these reports, sodium bicarbonate alone did not improve the pH or lactate concentrations, but bicarbonate-buffered peritoneal dialysis did. Perhaps importantly, peritoneal dialysis was very effective at removing lactate. The returned dialysate contained between 2.6 mEq/L and 14 mEq/L of lactate in one study,⁹⁰ and the calculated lactate clearance by peritoneal dialysis averaged 21 mL/min in another.⁹¹ The impact of bicarbonate infusion plus dialysis on cardiovascular function and outcome has not been studied systematically, nor has the relevance of the bicarbonate component of this potential treatment been examined.

CONCLUSION

Sodium bicarbonate is clearly effective in raising the arterial pH in critically ill patients with lactic acidosis. The impact on intracellular pH is unknown in such patients, but extrapolation from extensive

animal studies suggests that it is negative. Despite the correction of arterial acidemia, sodium bicarbonate, like DCA, has no favorable cardiovascular effects, even for patients with severe acidemia and receiving continuous infusions of catecholamines. Although hemodynamic improvement is not the only mechanism by which bicarbonate might be beneficial, animal studies have failed to yield alternatives. Even theoretical arguments in favor of sodium bicarbonate administration rely on a naïve representation of acid-base physiology, ignoring the complex compartmentalization of pH, the second-level effects of bicarbonate infusion, the impact of carbon dioxide generation, or the negative consequences of hyperlactatemia. We believe most clinicians who continue to use bicarbonate for patients with severe lactic acidosis do so largely because of their inclination to action: How can I “fail” to give bicarbonate when no alternative therapy is available and the mortality of this condition is so high?

The oft-cited rationale for bicarbonate use, that it might ameliorate the hemodynamic depression of metabolic acidemia, has been disproved convincingly. Any future role for bicarbonate in these patients depends on the formulation of new hypotheses of efficacy followed by animal and clinical studies to seek to confirm any proposed benefit. Given the current lack of evidence supporting its use, we cannot condone bicarbonate administration for patients with lactic acidosis. We extend this to those with pH < 7.2 on vasoactive drugs, inasmuch as bicarbonate has no measurable beneficial effects even in these sickest patients. Indeed, we do not give or advise bicarbonate infusion regardless of the pH.

REFERENCES

- 1 Weil MH, Afifi AA. Experimental and clinical studies on lactate and pyruvate as indicators of the severity of acute circulatory failure (shock). *Circulation* 1970; 41:989–1001
- 2 Cooper DJ, Walley KR, Wiggs BR, et al. Bicarbonate does not improve hemodynamics in critically ill patients who have lactic acidosis: a prospective, controlled clinical study. *Ann Intern Med* 1990; 112:492–498
- 3 Stacpoole PW, Wright EC, Baumgartner TG, et al. A controlled clinical trial of dichloroacetate for treatment of lactic acidosis in adults. *N Engl J Med* 1992; 327:1564–1569
- 4 Stacpoole PW, Wright EC, Baumgartner TG, et al. Natural history and course of acquired lactic acidosis in adults. *Am J Med* 1994; 97:47–54
- 5 Minot AS, Dodd K, Saunders JM. The acidosis of guanidine intoxication. *J Clin Invest* 1934; 13:917–932
- 6 Adrogue HJ, Madias NE. Management of life-threatening acid-base disorders: second of two parts. *N Engl J Med* 1998; 338:107–111
- 7 Oliva PB. Lactic acidosis. *Am J Med* 1970; 48:209–225
- 8 Warner A, Vaziri ND. Treatment of lactic acidosis. *South Med J* 1981; 74:841–847
- 9 Bonventre JV, Cheung JY. Effects of metabolic acidosis on

- viability of cells exposed to anoxia. *Am J Physiol* 1985; 249:C149–C159
- 10 Poole-Wilson PA, Langer GA. Effect of pH on ionic exchange and function in rat and rabbit myocardium. *Am J Physiol* 1975; 229:570–581
 - 11 Shapiro JI. Functional and metabolic responses of isolated hearts to acidosis: effects of sodium bicarbonate and Carbi-carb. *Am J Physiol* 1990; 258:H1835–H1839
 - 12 Cingolani HE, Faulkner SL, Mattiazzi AR, et al. Depression of human myocardial contractility with “respiratory” and “metabolic” acidosis. *Surgery* 1975; 77:427–432
 - 13 Nakanishi T, Okuda H, Kamata K, et al. Influence of acidosis on inotropic effect of catecholamines in newborn rabbit hearts. *Am J Physiol* 1987; 253:H1441–H1448
 - 14 Orchard CH, Kentish JC. Effects of pH change on the contractile function of cardiac muscle. *Am J Physiol* 1990; 258:C967–C981
 - 15 Downing SE, Talner NS, Gardner TH. Cardiovascular responses to metabolic acidosis. *Am J Physiol* 1965; 208:237–242
 - 16 Cooper DJ, Herbertson MJ, Werner HA, et al. Bicarbonate does not increase left ventricular contractility during L-lactic acidemia in pigs. *Am Rev Respir Dis* 1993; 148:317–322
 - 17 Wildenthal K, Mierziak DS, Myers RW, et al. Effects of acute lactic acidosis on left ventricular performance. *Am J Physiol* 1968; 214:1352–1359
 - 18 Thorens J-B, Jolliet P, Ritz M, et al. Effects of rapid permissive hypercapnia on hemodynamics, gas exchange, and oxygen transport and consumption during mechanical ventilation for the acute respiratory distress syndrome. *Intensive Care Med* 1996; 22:182–191
 - 19 Feihl F, Perret C. Permissive hypercapnia: how permissive should we be? *Am J Respir Crit Care Med* 1994; 150:1722–1737
 - 20 Potkin RT, Swenson ER. Resuscitation from severe acute hypercapnia: determinants of tolerance and survival. *Chest* 1992; 102:1742–1745
 - 21 Tuxen DV, Williams TJ, Scheinkestel CD, et al. Use of a measurement of pulmonary hyperinflation to control the level of mechanical ventilation in patients with acute severe asthma. *Am Rev Respir Dis* 1992; 146:1136–1142
 - 22 Hickling KG, Walsh J, Henderson S, et al. Low mortality rate in adult respiratory distress syndrome using low-volume, pressure-limited ventilation with permissive hypercapnia: a prospective study. *Crit Care Med* 1994; 22:1568–1578
 - 23 Bidani A, Tzouanakis AE, Cardenas VJ, et al. Permissive hypercapnia in acute respiratory failure. *JAMA* 1994; 272:957–962
 - 24 Kiely DG, Cargill RI, Lipworth BJ. Effects of hypercapnia on hemodynamic, inotropic, lusitropic, and electrophysiologic indices in humans. *Chest* 1996; 109:1215–1221
 - 25 Roupie E, Dambrosio M, Servillo G, et al. Titration of tidal volume and induced hypercapnia in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1995; 152:121–128
 - 26 Gamba G, Oseguera J, Castrejon M, et al. Bicarbonate therapy in severe diabetic ketoacidosis: a double-blind, randomized, placebo controlled trial. *Rev Invest Clin* 1991; 43:234–238
 - 27 Goldstein B, Shnnon DC, Todres ID. Supercarbia in children: clinical course and outcome. *Crit Care Med* 1990; 18:166–168
 - 28 Meldrum BS, Horton RW. Physiology of status epilepticus in primates. *Arch Neurol* 1973; 28:1–9
 - 29 Beresford HR, Posner JB, Plum F. Changes in brain lactate during induced cerebral seizures. *Arch Neurol* 1969; 20:243–248
 - 30 Glaser GH. Medical complications of status epilepticus. *Neurology* 1983; 34:395–397
 - 31 Marsh JD, Margolis TI, Kim D. Mechanism of diminished contractile response to catecholamines during acidosis. *Am J Physiol* 1988; 254:H20–H27
 - 32 Campbell GS, Houle DB, Crisp NW, et al. Depressed response to intravenous sympathomimetic agents in humans during acidosis. *Dis Chest* 1958; 33:18–22
 - 33 von Planta I, Weil MH, von Planta M, et al. Hypercarbic acidosis reduces cardiac resuscitability. *Crit Care Med* 1991; 19:1177–1182
 - 34 Kette F, Weil MH, von Planta M, et al. Buffer agents do not reverse intramyocardial acidosis during cardiac resuscitation. *Circulation* 1990; 81:1660–1666
 - 35 Kerber RE, Pandian NG, Hoyt R, et al. Effect of ischemia, hypertrophy, hypoxia, acidosis, and alkalosis on canine defibrillation. *Am J Physiol* 1983; 244:H825–H831
 - 36 Rose CE, Benthuyzen KV, Jackson JT, et al. Right ventricular performance during increased afterload impaired by hypercapnic acidosis in conscious dogs. *Circ Res* 1983; 52:76–84
 - 37 Bersentes TJ, Simmons DH. Effects of acute acidosis on renal hemodynamics. *Am J Physiol* 1967; 212:633–640
 - 38 Juan G, Calverley P, Talamo C, et al. Effect of carbon dioxide on diaphragmatic function in human beings. *N Engl J Med* 1984; 310:874–879
 - 39 Yanos J, Wood LDH, Davis K, et al. The effect of respiratory and lactic acidosis on diaphragm function. *Am Rev Respir Dis* 1993; 147:616–619
 - 40 Gores GJ, Nieminen A-L, Wray BE, et al. Intracellular pH during “chemical hypoxia” in cultured rat hepatocytes. *J Clin Invest* 1989; 83:386–396
 - 41 Preckel B, Schlack W, Obal D, et al. Effect of acidotic blood reperfusion on reperfusion injury after coronary artery occlusion in the dog heart. *J Cardiovasc Pharmacol* 1998; 31:179–186
 - 42 Kitakaze M, Takashima S, Funaya H, et al. Temporary acidosis during reperfusion limits myocardial infarct size in dogs. *Am J Physiol* 1997; 272:H2071–H2078
 - 43 Yatani A, Fujino T, Kinoshita K, et al. Excess lactate modulates ionic currents and tension components in frog atrial muscle. *J Mol Cell Cardiol* 1981; 13:147–161
 - 44 Stewart PA. How to understand acid-base: a quantitative acid-base primer for biology and medicine. New York, NY: Elsevier 1981; 1–179
 - 45 Sun JH, Filley GF, Hord K, et al. Carbi-carb: an effective substitute for NaHCO₃ for the treatment of acidosis. *Surgery* 1987; 5:835–839
 - 46 Arieff AI, Leach W, Park R, et al. Systemic effects of NaHCO₃ in experimental lactic acidosis in dogs. *Am J Physiol* 1982; 242:F586–F591
 - 47 Graf H, Leach W, Arieff AI. Metabolic effects of sodium bicarbonate in hypoxic lactic acidosis in dogs. *Am J Physiol* 1985; 249:F630–F635
 - 48 Benjamin E, Oropello JM, Abalos AM, et al. Effects of acid-base correction on hemodynamics, oxygen dynamics, and resuscitability in severe canine hemorrhagic shock. *Crit Care Med* 1994; 22:1616–1623
 - 49 Rhee KH, Toro LO, McDonald GG, et al. Carbi-carb, sodium bicarbonate, and sodium chloride in hypoxic lactic acidosis: effect on arterial blood gases, lactate concentrations, hemodynamic variables, and myocardial intracellular pH. *Chest* 1993; 104:913–918
 - 50 Fraley DS, Adler S, Bruns FJ, et al. Stimulation of lactate production by administration of bicarbonate in a patient with a solid neoplasm and lactic acidosis. *N Engl J Med* 1980; 303:1100–1102
 - 51 Bellingham AJ, Detter JC, Lenfant C. Regulatory mecha-

- nisms of hemoglobin oxygen affinity in acidosis and alkalosis. *J Clin Invest* 1971; 50:700–706
- 52 Park R, Arieff AI. Treatment of lactic acidosis with dichloroacetate in dogs. *J Clin Invest* 1982; 70:853–862
 - 53 Beech JS, Nolan KM, Iles RA, et al. The effects of sodium bicarbonate and a mixture of sodium bicarbonate and carbonate (Carbicarb) on skeletal muscle pH and hemodynamic status in rats with hypovolemic shock. *Metabolism* 1994; 43:518–522
 - 54 Klepper ID, Kucera RF, Kindig NB, et al. A comparative study of sodium bicarbonate and Carbicarb in the treatment of metabolic acidosis induced by hemorrhagic shock. *J Crit Care* 1988; 3:256–261
 - 55 Beech JS, Williams SCR, Iles RA, et al. Hemodynamic and metabolic effects in diabetic ketoacidosis in rats of treatment with sodium bicarbonate or a mixture of sodium bicarbonate and sodium carbonate. *Diabetologia* 1995; 38:889–898
 - 56 Mathieu D, Nevier R, Billard V, et al. Effects of bicarbonate therapy on hemodynamics and tissue oxygenation in patients with lactic acidosis: a prospective, controlled clinical study. *Crit Care Med* 1991; 19:1352–1356
 - 57 Bureau MA, Begin R, Berthiaume Y, et al. Cerebral hypoxia from bicarbonate infusion in diabetic acidosis. *J Pediatr* 1980; 96:968–973
 - 58 Bishop RL, Weisfeldt ML. Sodium bicarbonate administration during cardiac arrest: effect on arterial pH, PCO₂, and osmolality. *JAMA* 1976; 235:506–509
 - 59 Posner JB, Plum F. Spinal-fluid pH and neurologic symptoms in systemic acidosis. *N Engl J Med* 1967; 277:605–613
 - 60 Bollaert PE, Robin-Lherbier B, Mallie JP, et al. Effects of sodium bicarbonate on striated muscle metabolism and intracellular pH during endotoxic shock. *Shock* 1994; 1:196–200
 - 61 Thompson CH, Syme PD, Williams EM, et al. Effect of bicarbonate administration on skeletal muscle intracellular pH in the rat: implications for acute administration of bicarbonate in man. *Clin Sci* 1992; 82:559–564
 - 62 Bersin RM, Arieff AI. Improved hemodynamic function during hypoxia with Carbicarb, a new agent for the management of acidosis. *Circulation* 1988; 77:227–233
 - 63 Shapiro JI, Whalen M, Kucera R, et al. Brain pH responses to sodium bicarbonate and Carbicarb during systemic acidosis. *Am J Physiol* 1989; 256:H1316–H1321
 - 64 Shapiro JI, Whalen M, Chan L. Hemodynamic and hepatic pH responses to sodium bicarbonate and Carbicarb during systemic acidosis. *Magn Reson Med* 1990; 16:403–410
 - 65 Nakashima K, Yamashita T, Kashiwagi S, et al. The effect of sodium bicarbonate on CBF and intracellular pH in man: stable Xe-CT and ³¹P-MRS. *Acta Neurol Scand Suppl* 1996; 166:96–98
 - 66 Bjerneroth G, Sammeli O, Li YC, et al. Effects of alkaline buffers on cytoplasmic pH in lymphocytes. *Crit Care Med* 1994; 22:1550–1556
 - 67 Ritter JM, Doktor HS, Benjamin N. Paradoxical effect of bicarbonate on cytoplasmic pH. *Lancet* 1990; 335:1243–1246
 - 68 Arieff AI, Leach WJ, Lazarowitz VC. Effects of NaHCO₃ in therapy of experimental lactic acidosis [abstract]. *Kidney Int* 1978; 14:645–645
 - 69 Goldsmith DJA, Forni LG, Hilton PJ. Bicarbonate therapy and intracellular acidosis. *Clin Sci* 1997; 93:593–598
 - 70 Levraut J, Labib Y, Chave S, et al. Effect of sodium bicarbonate on intracellular pH under different buffering conditions. *Kidney Int* 1996; 49:1262–1267
 - 71 Sirieix D, DeLayance S, Paris M, et al. Tris-hydroxymethyl aminomethane and sodium bicarbonate to buffer metabolic acidosis in an isolated heart model. *Am J Respir Crit Care Med* 1997; 155:957–963
 - 72 Tanaka M, Nishikawa T, Mizutani T. Normovolemic hemodilution attenuates cardiac depression induced by sodium bicarbonate in canine metabolic acidosis. *Br J Anaesth* 1996; 77:408–412
 - 73 Basir MA, Bhatia J, Brudno DS, et al. Effects of Carbicarb and sodium bicarbonate on hypoxic lactic acidosis in newborn pigs. *J Invest Med* 1996; 44:70–74
 - 74 Tanaka M, Nishikawa T. Acute hemodynamic effects of sodium bicarbonate administration in respiratory and metabolic acidosis in anesthetized dogs. *Anaesth Intensive Care* 1997; 25:615–620
 - 75 Lever E, Jaspan JB. Sodium bicarbonate therapy in severe diabetic ketoacidosis. *Am J Med* 1983; 75:263–268
 - 76 Morris LR, Murphy MB, Kitabchi AE. Bicarbonate therapy in severe diabetic ketoacidosis. *Ann Intern Med* 1986; 105:836–840
 - 77 Hale PJ, Crase J, Natrass M. Metabolic effects of bicarbonate in the treatment of diabetic ketoacidosis. *BMJ* 1984; 289:1035–1038
 - 78 Okuda Y, Adrogué HJ, Field JB, et al. Counterproductive effects of sodium bicarbonate in diabetic ketoacidosis. *J Clin Endocrinol Metab* 1996; 81:314–320
 - 79 Vukmir RB, Bircher NG, Radovsky A, et al. Sodium bicarbonate may improve outcome in dogs with brief or prolonged cardiac arrest. *Crit Care Med* 1995; 23:515–522
 - 80 Kette F, Weil MH, Gazmuri RJ. Buffer solutions may compromise cardiac resuscitation by reducing coronary perfusion pressure. *JAMA* 1991; 266:2121–2126
 - 81 Roberts D, Landolfo K, Light RB, et al. Early predictors of mortality for hospitalized patients suffering cardiopulmonary arrest. *Chest* 1990; 97:413–419
 - 82 Máttar JA, Weil MH, Shubin H, et al. Cardiac arrest in the critically ill: II. Hyperosmolal states following cardiac arrest. *Am J Med* 1974; 56:162–168
 - 83 Huseby JS, Gumprecht DG. Hemodynamic effects of rapid bolus hypertonic sodium bicarbonate. *Chest* 1981; 79:552–554
 - 84 Bersin RM, Chatterjee K, Arieff AI. Metabolic and hemodynamic consequences of sodium bicarbonate administration in patients with heart disease. *Am J Med* 1989; 87:7–14
 - 85 Fields ALA, Wolman SL, Halperin ML. Chronic lactic acidosis in a patient with cancer: therapy and metabolic consequences. *Cancer* 1981; 47:2026–2029
 - 86 Lang RM, Fellner SK, Neumann A, et al. Left ventricular contractility varies directly with blood ionized calcium. *Ann Intern Med* 1988; 108:524–529
 - 87 Stacpoole PW. The pharmacology of dichloroacetate. *Metabolism* 1989; 38:1124–1144
 - 88 Stacpoole PW, Lorenz AC, Thomas RG, et al. Dichloroacetate in the treatment of lactic acidosis. *Ann Intern Med* 1988; 108:58–63
 - 89 Stacpoole PW, Harman EM, Curry SH, et al. Treatment of lactic acidosis with dichloroacetate. *N Engl J Med* 1983; 309:390–399
 - 90 Vaziri ND, Ness R, Wellikson L, et al. Bicarbonate-buffered peritoneal dialysis: an effective adjunct in the treatment of lactic acidosis. *Am J Med* 1979; 67:392–396
 - 91 Sheppard JM, Lawrence JR, Oon RCS, et al. Lactic acidosis: recovery associated with use of peritoneal dialysis. *Aust N Z J Med* 1972; 4:389–392